



HAL
open science

INHIBITION OF THE FUNGAL PATHOGEN METARHIZIUM ANISOPLIAE IN THE ALIMENTARY TRACTS OF FIVE TERMITE (ISOPTERA) SPECIES

Thomas Chouvenc, Nan-Yao Su, Alain Robert

► **To cite this version:**

Thomas Chouvenc, Nan-Yao Su, Alain Robert. INHIBITION OF THE FUNGAL PATHOGEN METARHIZIUM ANISOPLIAE IN THE ALIMENTARY TRACTS OF FIVE TERMITE (ISOPTERA) SPECIES. Florida Entomologist, 2010, 93 (3), pp.467-469. 10.1653/024.093.0327 . bioemco-00526603

HAL Id: bioemco-00526603

<https://hal-bioemco.ccsd.cnrs.fr/bioemco-00526603v1>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Inhibition of the Fungal Pathogen *Metarhizium anisopliae* in the Alimentary Tracts of Five Termite (Isoptera) Species

Author(s): Thomas Chouvenc, Nan-Yao Su and Alain Robert

Source: Florida Entomologist, 93(3):467-469.

Published By: Florida Entomological Society

DOI: <http://dx.doi.org/10.1653/024.093.0327>

URL: <http://www.bioone.org/doi/full/10.1653/024.093.0327>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

INHIBITION OF THE FUNGAL PATHOGEN *METARHIZIUM ANISOPLIAE* IN THE ALIMENTARY TRACTS OF FIVE TERMITE (ISOPTERA) SPECIESTHOMAS CHOUVENC^{1*}, NAN-YAO SU¹ AND ALAIN ROBERT^{2,3}¹Department of Entomology and Nematology, Ft. Lauderdale Research and Education Center, University of Florida, Institute of Food and Agricultural Sciences, 3205 College Ave, Ft. Lauderdale, FL 33314, USA²UMR 5548 Communication chimique et développement chez les insectes, Université de Bourgogne, 6 Bd Gabriel, 2100 Dijon, France³Current address: Centre IRD France Nord, 32 avenue Varagnat, 93143 Bondy, France

*Corresponding author

Supplemental material online at [http://www.fcla.edu/FlaEnt/fe93\(3\).htm#InfoLink1](http://www.fcla.edu/FlaEnt/fe93(3).htm#InfoLink1)

There are more than 2,600 described species of termites, from 7 families, and they are adapted to a wide range of habitats resulting in high diversity in morphology, physiology, behavior, nesting ecology, and associated microbial community (Abe et al. 2000). Sociality is thought to be a key factor in their ecological success and diversity, but how termites evolved to resist the increased pressure from pathogens as they moved from the solitary lifestyle of the wood roach-like ancestor towards eusociality remains poorly understood (Fefferman et al. 2007). Chouvenc and Su (2010) showed that the subterranean termite *Reticulitermes flavipes* (Kollar) has the capacity to prevent epizootics of the entomopathogenic fungus *Metarhizium anisopliae* (Metsch.) Sorokin due to synergy among defense mechanisms. Chouvenc et al. (2009a) described *in vivo* the antifungal activity of the alimentary tract in *R. flavipes* and showed that after grooming and ingestion of conidia of *M. anisopliae*, germination did not occur in the alimentary tract of living termites. In dead termites, this antifungal activity remained strong at least 2 d *post mortem* throughout all gut regions.

We hypothesized that gut antifungal activity is an ancestral feature and was maintained during the radiation of the isopteran lineage. To test this, we extended our observations to determine if this mechanism exists in 5 additional termite species from 5 different families. Histological preparation was identical to Chouvenc et al. (2009a) and was performed on selected specimens described in Chouvenc et al. (2009c). Histological analysis included 25 *Hodotermopsis sjoestedti* Holmgren (Termopsidae), 8 *Hodotermes mossambicus* (Hagen) (Hodotermitidae), 13 *Kaloterms flavicollis* (Fabricius) (Kalotermitidae), 10 *Prorhinotermes canalifrons* (Sjöstedt) (Rhinotermitidae), and 2 *Nasutitermes voeltzkowi* (Wasmann) (Termitidae). The presence or absence of conidia in the gut was recorded in all specimens, and the presence of fungal hyphae was examined in healthy, moribund, and dead specimens. Full histological observations for each species are available in the supplemental information.

After exposure to a solution of *M. anisopliae* conidia, grooming was performed by the termites and conidia were subsequently found in the alimentary tracts of the groomers for all 5 termite species (Fig. 1A). For each species, conidia in the alimentary tract did not germinate, as previously described in *R. flavipes* (Chouvenc et al. 2009a). In all the specimens that were fixed after their death (usually 2-5 d after fungal exposure) for histological preparation, the fungal hyphae quickly spread in the hemocoel, the muscles (Fig. 1B) and the fat body within the 24 h *post mortem*, but did not readily invade the salivary glands. The fungus did not penetrate the salivary glands until at least after 1 d *post mortem*. In addition, *M. anisopliae* hyphae did not penetrate the gut lumen until at least after 2 d *post mortem*. We confirmed that the fungistatic activity of the alimentary tract was present in all tested species and that this activity remained strong in the alimentary tract at least 2 d *post mortem* (Figs. 1C and 1D), although the salivary glands were already invaded by the fungus at this time (Fig. 1E). Therefore, all conidia found in the digestive tract of live termites did not germinate and the fungistatic activity of the gut in all termite species remained active for several days after the death of the termites, as previously showed in *R. flavipes* (Chouvenc et al. 2009a). We suggest that gut fungistatic activity is an ancestral trait that has been conserved during the radiation of the Isoptera. However, we previously showed that all termites studied had a variable mortality rate when exposed to *M. anisopliae* (Chouvenc et al. 2009c). The present results suggest that this difference of susceptibility among species may depend on other physiological factors, such as the cellular immunity (Chouvenc et al. 2009b), or other behavioral and physiological factors (Cremier et al. 2007). It has been suggested that part of this fungistatic activity is due to the production of antimicrobial peptides in the salivary glands (Lamberty et al. 2001; Bulmer & Crozier 2004), but our results showed that in all species the salivary glands were invaded by the fungus much




Fig. 1. Observation of *Metarhizium anisopliae* in termites alimentary tract of 6 species after histological preparation. (A) *Nasutitermes voeltzkowi*, ungerminated conidium in cuticular foldings of the crop (foregut). (B) *Hodotermes mossambicus*, growth of hyphae in the hemocoel and in the muscles at 1 d *post mortem*. (C) *Hodotermopsis sjoestedti*, growth of hyphal bodies in the muscles around the midgut but absence of penetration of the midgut lumen by the fungus at 1 d *post mortem*. (D) *Hodotermopsis sjoestedti*, beginning of penetration by the fungus of the hindgut lumen at 3 d *post mortem*. (E) *Prorehinotermes canalifrons*, invasion of the salivary glands at 2 d *post mortem*. (F) *Kaloterмес flavicollis*, fragment of a cellular encapsulation from the external cuticle of a cannibalized nest-mate in the midgut. *co* = conidia, *ct* = external cuticle, *fb* = fat body, *fgc* = foregut cuticle, *hgc* = hindgut cuticle, *hy* = hyphae, *mel* = melanization, *mep* = midgut epithelium, *mt* = malpighian tubule, *ms* = muscle, *pt* = protozoan, *sg* = salivary gland, * = gut lumen. The scale bars represent 50 μm .

faster than the gut lumen, indicating that the antimicrobial compounds contained in the salivary glands may have a limited activity inside the digestive tract, or that the fungistatic activity of the gut is due to a range of various compounds.

Finally, we observed in the midgut of a *K. flavicollis* specimen the presence of a fragment of external cuticle with a cellular encapsulation (Fig. 1F), as described by Chouvenec et al. (2009b). This observation suggests that the termite performed

necrophagy on a dead nestmate that has been previously infected by *M. anisopliae*, and that the passage through the gut lumen did not allow this ingested fungal particles to grow while in the alimentary tract. Therefore, cannibalism of an infected cadaver may aid in preventing *M. anisopliae* from completing its life cycle in a termite group. Grooming and necrophagy in association with gut antifungal activity provides an environment that reduces the chances for the fungus to produce an epizootic into the termite nest. This mechanism appeared to be conserved in all tested species from 5 different families.

SUMMARY

We previously showed that in the subterranean termite *Reticulitermes flavipes* conidia of *Metarhizium anisopliae* did not germinate in the alimentary tract of living termites and the antifungal activity remained for 2 d *post mortem* in the gut. We have confirmed that this mechanism is common throughout the termite phylogeny by extending our observations to 5 additional species, from 5 different families. Grooming and ingestion of microorganisms from the cuticle of nestmate was found to be consistent in all the species and none of the conidia found in the alimentary tracts germinated.

REFERENCES CITED

- ABE, T. 1987. Evolution of life types in termites, pp. 125-148. In S. Kawano, J. H. Connell, and T. Hidaka (eds.), *Evolution and Coadaptation: Biotic Communities*. University of Tokyo Press, Tokyo.
- BULMER, M., AND CROZIER, R. H. 2004. Duplication and diversifying selection among termite antifungal peptides. *Mol. Biol. Evol.* 21: 2256-2264.
- CHOUVENC, T., AND SU, N.-Y. 2010. Apparent synergy among defense mechanisms in subterranean termites (Rhinotermitidae) against epizootic events - The limits and potential for biological control. *J. Econ. Entomol.* In press.
- CHOUVENC, T., SU, N.-Y., AND ROBERT, A. 2009a. Inhibition of *Metarhizium anisopliae* in the alimentary tract of the eastern subterranean termite *Reticulitermes flavipes*. *J. Invertebr. Pathol.* 101: 130-136.
- CHOUVENC, T., SU, N.-Y., AND ROBERT, A. 2009b. Cellular encapsulation in the eastern subterranean termite, *Reticulitermes flavipes* (Isoptera), against infection by the entomopathogenic fungus *Metarhizium anisopliae*. *J. Invertebr. Pathol.* 101: 234-241.
- CHOUVENC, T., SU, N.-Y., AND ROBERT, A. 2009c. Susceptibility of seven termite species (Isoptera) to the entomopathogenic fungus *Metarhizium anisopliae*. *Sociobiology* 54, 723-748.
- CREMER, S., ARMITAGE, S. A. O., AND SCHMID-HEMPEL, P. 2007. Social immunity. *Curr. Biol.* 17: 693-702.
- CULLINEY, T. W., AND GRACE, J. K. 2000. Prospects for the biological control of subterranean termites (Isoptera: Rhinotermitidae), with special reference to *Coptotermes formosanus*. *Bull. Entomol. Res.* 90: 9-21.
- FEFFERMAN, N. H., TRANIELLO, J. F. A., ROSENGAUS, R. B., AND CALLERI II, D. V. 2007. Disease prevention and resistance in social insects: modeling the survival consequences of immunity, hygienic behavior, and colony organization. *Behav. Ecol. Sociobiol.* 61, 565-577.
- LAMBERTY, M., ZACHARY, D., LANOT, R., BORDEREAU, C., ROBERT, A., HOFFMANN, J. A., AND BULET, P. 2001. Insect immunity constitutive expression of a cysteine-rich antifungal and a linear antibacterial peptide in a termite insect. *J. Biol. Chem.* 276: 4085-4092.